

Drug Facts

Active ingredient ***Purpose***
(in each tablet)

Magnesium hydroxide 400 mg.....Saline laxative

Uses

- for relief of occasional constipation
 - this product usually produces bowel movement in 1/2 to 6 hours
-

Warnings

Ask a doctor before using any laxative if your child has

- kidney disease
 - a magnesium-restricted diet
 - abdominal pain, nausea, or vomiting
 - a sudden change in bowel habits lasting more than 2 weeks
 - constipation that continues after 1 week of use, contact your child's doctor
 - already used a laxative for more than 1 week
-

Stop using this product and consult a doctor if your child

- has rectal bleeding
- does not have a bowel movement within 6 hours of taking this product

These symptoms may be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.



Drug Facts (continued)

Directions

Use dosage chart for proper dosing. Doses may be taken as a single daily dose or in divided doses. Drink a full glass (8 ounces) of liquid with each dose.

Age	Starting Dose	Maximum dose per day (24 hours)
children 6 to under 12 years	3-6 tablets	6 tablets
children 2 to under 6 years	1-3 tablets	3 tablets
children under 2 years	ask a doctor	

Other information

- each tablet contains: magnesium 170 mg
- CHILD RESISTANT CAP
- The top of the bottle is sealed with foil for your safety. Do not use if foil imprinted "SEALED for YOUR PROTECTION" is broken or missing.
- Gluten Free & Sugar Free

Inactive ingredients

colloidal silicon dioxide, FD&C red #40 aluminum lake, flavor, magnesium stearate, maltodextrin, mannitol, sorbitol, stearic acid, sucralose

Questions? 1-866-255-6960 or www.pedia-lax.com