

Drug Facts

Active ingredient (in each 15 mL)	Purpose
Docusate Sodium 50 mg	Stool softener

Uses ■ for prevention of dry, hard stools ■ for relief of occasional constipation ■ this product usually produces a bowel movement in 12 to 72 hours

Warnings

Drug Interaction Precaution: Do not give this product to your child if your child is presently taking mineral oil, unless directed by a doctor.

Do not use when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before using any laxative if your child has

- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week
- if constipation continues after 1 week of use, contact your child's doctor

Stop using this product and consult a doctor if your child has

- rectal bleeding
 - no bowel movement within 72 hours of using this product
- These symptoms may be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center (1-800-222-1222) right away.

Directions

- do not give more than 45 mL in any 24 hour period
- use only with enclosed dosing cup
- find right dose on chart below
- mL = milliliter

Children 2 to under 12 years	15 mL to 45 mL; do not take more than 45 mL in 24 hours
Children under 2 years	ask a doctor

Doses can be diluted in a suitable beverage to prevent throat irritation and to help mask the naturally tart flavor of the active ingredient (see side panel for info).

Other information

- each 15 mL contains: **sodium 13 mg**
- child resistant cap
- the top of the bottle is sealed with foil for your safety. Do not use if foil imprinted "sealed for your protection" is broken or missing.
- gluten free

Inactive ingredients citric acid, edetate disodium, flavor, methylparaben, polyethylene glycol, povidone, propylene glycol, propylparaben, sodium citrate, sorbitol, sucralose, water, xanthan gum, xylitol

Questions? 1-866-255-6960 www.pedia-lax.com